

STAT 450: Statistical Theory

Statistical Inference

Definition: A **model** is a family $\{f_\theta; \theta \in \Theta\}$ of possible densities for some random variable X .

- Our data set is X , so X will generally be a big vector or matrix or even more complicated object.
- Can also specify models by things like cdf instead of density.
- We assume in this course: true density f of X is some f_{θ_0} for some $\theta_0 \in \Theta$.
- We call θ_0 the true value of the parameter.

- In real world this assumption will be wrong; we hope it is not wrong in an important way.
- If we are very worried that it is wrong we enlarge our model putting in more distributions and making Θ bigger.
- Theorists also try to describe the effects of this wrong assumption; subject is called *model mis-specification*.

- Goal: observe value of X , then guess θ_0 or some property of θ_0 .
- Specific classic mathematical versions of this:
 1. Point estimation: compute estimate $\hat{\theta} = \hat{\theta}(X)$ which lies in Θ (or something close to Θ).
 2. Point estimation of a function of θ : compute an estimate $\hat{\phi} = \hat{\phi}(X)$ of $\phi = g(\theta)$.
 3. Interval (or set) estimation: compute a set $C = C(X)$ in Θ which we think will contain θ_0 .
 4. Hypothesis testing: choose between $\theta_0 \in \Theta_0$ and $\theta_0 \notin \Theta_0$ where $\Theta_0 \subset \Theta$.
 5. Prediction: guess the value of an observable random variable Y whose distribution depends on θ_0 . Typically Y is the value of the variable X in a repetition of the experiment.

Statistical Philosophies

Several schools of statistical thinking.

Main schools of thought summarized roughly as follows:

- **Neyman Pearson:** Statistical procedure is evaluated by its long run frequency performance.

Imagine repeating the data collection exercise many times, independently.

Quality of procedure measured by its average performance when true distribution of X values is P_{θ_0} .

- **Bayes:** Treat θ as random just like X .
- Compute conditional law of unknown quantities given knowns.
- Ask how procedure will work on the data we actually got – no averaging over data we might have got.
- **Likelihood:** Try to combine previous 2 by looking only at actual data while trying to avoid treating θ as random.

We use Neyman Pearson approach to evaluate quality of likelihood and other methods.